

Name: MARK EGLY
Age: 61
Gender: Male

Bar Code: 0022080976
Control Number: 39594250

Test Date: Mar 09, 2017 02:01 PM
Last Food/Drink: Mar 02, 2017 07:21 PM
Specimen Collected: Mar 03, 2017 11:21 AM

Heart & Arteries

Test Name	Your Result	Units	Reference Range	Test Guide
Cholesterol	146	mg/dL	140 - 250	Cholesterol is a waxy, fat-like substance that occurs naturally in all parts of the body. Your body needs some cholesterol to work properly, but if you have too much in your blood, it can stick to the walls of your arteries. This is called plaque. Plaque can narrow your arteries or even block them.
HDL	58	mg/dL	35 - 80	High-Density Lipoprotein (HDL) is the "good" or "healthy" cholesterol, which helps keep the Low-Density Lipoprotein (LDL) "bad" cholesterol from getting lodged into your artery walls. A healthy level of HDL (greater than 40 mg/dL for men and greater than 50 mg/dL for women) may also protect against heart attack and stroke, while low levels of HDL have been shown to increase the risk of heart disease.
LDL	79	mg/dL	0 - 129	Low-Density Lipoprotein (LDL) cholesterol is the "bad" or "lousy" cholesterol. When too much of it circulates in the blood, it can clog arteries, increasing your risk of heart attack and stroke.
Cholesterol/HDL Ratio	2.5		0 - 6.5	The ratio of total Cholesterol to HDL-Cholesterol is another indicator of heart disease risk. A ratio is just one value divided by another number, but in medicine it can be helpful in predicting disease. A ratio of 5.0 or less is associated with a lower risk of heart disease.
LDL/HDL Ratio	1.36		0.9 - 5.3	The LDL/HDL Ratio may be reported as part of a lipid profile, a group of tests that are often ordered together to determine risk of heart disease and an important part of heart risk assessments. LDL/HDL Cholesterol Ratio is a calculated value that is an indicator of heart disease risk. The lower the ratio, the lower the risk.
Triglycerides	45	mg/dL	0 - 150	Triglycerides are fats (lipids) that provide a reserve of energy. Increases in triglycerides may indicate heart disease risk. Triglycerides can rise with obesity, diabetes and alcohol consumption.

Test Name	Your Result	Units	Reference Range	Test Guide
Diuretic SCRN- U	NEGATIVE	ng/mL	0 - 1499	A diuretic is any drug (often prescribed for high blood pressure) that elevates the frequency and amount of urination. This test detects the presence of a diuretic in urine.

Kidney & Bladder

Test Name	Your Result	Units	Reference Range	Test Guide
BUN	19	mg/dL	9 - 27	Blood urea nitrogen (BUN) is an end product of protein metabolism. The BUN test is primarily used, along with the creatinine test, to evaluate the kidney and to help diagnose kidney disease. It also may be used to evaluate a person's general health status.
Creatinine	1.1	mg/dL	0.7 - 1.5	Creatinine is a metabolic product released from muscle tissue and excreted from the kidneys. The test is used along with a Blood Urea Nitrogen (BUN) test to assess kidney function.
Urine PH Screen	5.5		4.0 - 8.7	This test measures how acidic the urine is.
Protein	6	mg/dL	0 - 30	A Protein urine test measures the amount of Proteins, such as Albumin, found in a urine sample. Urinary Protein elevations may indicate the presence of kidney disease, but levels vary with urine concentration.
Leukocyte Screen	NEGATIVE		NEGATIVE	Leukocyte Esterase is an enzyme in white blood cells. When present it may indicate infection of the kidney or urinary tract, including the bladder.
Hemoglobin Screen	NEGATIVE		NEGATIVE	Hemoglobin is a molecule attached to red blood cells that helps move oxygen and carbon dioxide through the body. Hemoglobin in the urine may indicate kidney and/or urinary tract disease but may also occur in normal conditions such as during menstruation.
Urine Creatinine	189.1	mg/dL	27.0 - 260.0	Creatinine is a breakdown product of Creatine, which is an important part of muscle. Creatinine is removed from the body entirely by the kidneys.
Protein/Creatinine Ratio	0.03	mg/mg Creat	0.00 - 0.20	Urine protein testing is used to detect protein in the urine. Creatinine is a breakdown product of creatine, which is an important part of muscle. This test measures creatinine with protein to calculate a urine protein/creatinine ratio (UP/CR). The test is used to evaluate kidney function as well as to detect other urinary tract disorders.

Liver

Test Name	Your Result	Units	Reference Range	Test Guide
Alkaline Phosphatase	48	U/L	30 - 125	Alkaline Phosphatase is an enzyme found primarily in the liver and bones. Elevated levels may indicate the presence of bone or liver disorders. The enzyme activity is also increased following fractures and in growing children and pregnant women.
Total Bilirubin	0.9	mg/dL	0.2 - 1.5	Bilirubin is a breakdown product of red blood cells. Abnormally high total bilirubin levels may occur in individuals with liver and gallbladder disease and may cause jaundice (yellowing of skin and eyes).
AST	27	U/L	0 - 41	Aspartate Aminotransferase (AST) is an enzyme found in the liver and in cardiac and skeletal muscle. AST may rise in liver, heart and muscle disorders but may also increase following strenuous, prolonged exercise.
ALT	26	U/L	0 - 45	Alanine Aminotransferase (ALT) is an enzyme found in the liver and rises with liver disease.
GGT	9	U/L	0 - 65	Gamma Glutamyl Transpeptidase (GGT) is a liver enzyme. It may rise with heavy alcohol consumption, certain medications and liver diseases.
Total Protein	6.8	g/dL	6.1 - 7.9	Total Protein measurements can reflect nutritional status and may be used to screen for and help diagnose kidney disease, liver disease and many other conditions. Protein in the blood includes two major components, Albumin and Globulin.
Albumin	4.2	g/dL	3.6 - 4.9	Albumin is the largest portion of total blood Protein. Decreased blood albumin may indicate many disorders, including poor nutrition and advanced liver disease.
Globulin	2.6	g/dL	2.1 - 3.6	Globulin is a major component of blood proteins. Abnormal levels (both elevated and decreased) may indicate infections, allergic states, immune disorders and other diseases.

Other

Test Name	Your Result	Units	Reference Range	Test Guide
Serum HIV	NEGATIVE		NEGATIVE	HIV antibody testing detects infection with the virus that causes AIDS. The test may not become positive until several weeks after exposure to the virus.

Test Name	Your Result	Units	Reference Range	Test Guide
PSA	0.82	ng/mL	0.00 - 4.00	Prostate-Specific Antigen (PSA) is a test in males that can detect prostate abnormalities including cancer and an enlarged prostate. Early detection and follow-up with your physician is the key to successful treatment and reduced mortality.

Cotinine	NEGATIVE	mcg/mL	0 - 0.29	Cotinine is the major by-product of nicotine. Its presence indicates tobacco use.
----------	----------	--------	----------	---

Pancreas

Test Name	Your Result	Units	Reference Range	Test Guide
Urine Glucose	NEGATIVE	g/dL	0.00 - 0.24	The Urine Glucose test measures the amount of glucose (sugar) in a urine sample. The presence of glucose in the urine is called glycosuria or glucosuria. Glucose is not normally present in urine, but may occur in diabetes and other illnesses.

Physical Measurements Performed By Paramedical Examiner

Height - standard: 6' 3"	Blood Pressure Reading 1: 110/66
Height - metric: 190.5 cm	Blood Pressure Reading 2: 110/60
Weight - standard: 168 lbs.	Blood Pressure Reading 3: 108/60
Weight - metric: 76.2 kg	Pulse Reading 1: 65
BMI: 20.9	Pulse Reading 2: 0



Welcome, MARK English Español Français | My Account | Log Out

Ocular List

Physical Measurements View/Print Results

MARK EGLY's Laboratory Results

Sort By: **Default** | In Range | Out of Range

- Heart & Arteries
- Kidney & Bladder
- Liver
- Pancreas
- Other
- All**

Cholesterol Heart & Arteries

Cholesterol is a waxy, fat-like substance that occurs naturally in all parts of the body. Your body needs some cholesterol to work properly, but if you have too much in your blood, it can stick to the walls of your arteries. This is called plaque. Plaque can narrow your arteries or even block them.

Your Result

146
mg/dL



Reference Range: 140 - 250

HDL Heart & Arteries

High-Density Lipoprotein (HDL) is the "good" or "healthy" cholesterol, which helps keep the Low-Density Lipoprotein (LDL) "bad" cholesterol from getting lodged into your artery walls. A healthy level of HDL (greater than 40 mg/dL for men and greater than 50 mg/dL for women) may also protect against heart attack and stroke, while low levels of HDL have been shown to increase the risk of heart disease.

Your Result

58
mg/dL



Reference Range: 35 - 80

LDL Heart & Arteries

Low-Density Lipoprotein (LDL) cholesterol is the "bad" or "lousy" cholesterol. When too much of it circulates in the blood, it can clog arteries, increasing your risk of heart attack and stroke.

Your Result

79
mg/dL



Reference Range: 0 - 129

Cholesterol/HDL Ratio Heart & Arteries

The ratio of total Cholesterol to HDL-Cholesterol is another indicator of heart disease risk. A ratio is just one value divided by another number, but in medicine it can be helpful in predicting disease. A ratio of 5.0 or less is associated with a lower risk of heart disease.

Your Result

2.5



Reference Range: 0 - 6.5

LDL/HDL Ratio Heart & Arteries

The LDL/HDL Ratio may be reported as part of a lipid profile, a group of tests that are often ordered together to determine risk of heart disease and an important part of heart risk assessments. LDL/HDL Cholesterol Ratio is a calculated value that is an indicator of heart disease risk. The lower the ratio, the lower the risk.

Your Result

1.36



Reference Range: 0.9 - 5.3

Triglycerides Heart & Arteries

Triglycerides are fats (lipids) that provide a reserve of energy. Increases in triglycerides may indicate heart disease risk. Triglycerides can rise with obesity, diabetes and alcohol consumption.

Your Result

45
mg/dL

▲
Reference Range: 0 - 150

Diuretic SCRN- U Heart & Arteries

A diuretic is any drug (often prescribed for high blood pressure) that elevates the frequency and amount of urination. This test detects the presence of a diuretic in urine.

Your Result

NEGATIVE

BUN Kidney & Bladder

Blood urea nitrogen (BUN) is an end product of protein metabolism. The BUN test is primarily used, along with the creatinine test, to evaluate the kidney and to help diagnose kidney disease. It also may be used to evaluate a person's general health status.

Your Result

19
mg/dL

▲
Reference Range: 9 - 27

Creatinine Kidney & Bladder

Creatinine is a metabolic product released from muscle tissue and excreted from the kidneys. The test is used along with a Blood Urea Nitrogen (BUN) test to assess kidney function.

Your Result

1.1
mg/dL

▲
Reference Range: 0.7 - 1.5

Urine PH Screen Kidney & Bladder

This test measures how acidic the urine is.

Your Result

5.5

▲
Reference Range: 4.0 - 8.7

Protein Kidney & Bladder

A Protein urine test measures the amount of Proteins, such as Albumin, found in a urine sample. Urinary Protein elevations may indicate the presence of kidney disease, but levels vary with urine concentration.

Your Result

6
mg/dL

▲
Reference Range: 0 - 30